

Bachman's™

Growing Since 1885



New England Aster

Aster novae-angliae

Height: 3 feet

Spread: 24 inches

Sunlight: ☉ ●

Hardiness Zone: 4

Other Names: New England Aster, Michaelmas Daisy

Ornamental Features

New England Aster has masses of beautiful pink daisy flowers with yellow eyes at the ends of the stems from early to late fall, which are most effective when planted in groupings. The flowers are excellent for cutting. Its narrow leaves remain dark green in color throughout the season. The fruit is not ornamentally significant.

Landscape Attributes

New England Aster is a dense herbaceous perennial with a mounded form. Its relatively fine texture sets it apart from other garden plants with less refined foliage.

This is a relatively low maintenance plant, and is best cleaned up in early spring before it resumes active growth for the season. It is a good choice for attracting butterflies to your yard, but is not particularly attractive to deer who tend to leave it alone in favor of tastier treats. Gardeners should be aware of the following characteristic(s) that may warrant special consideration;

- Disease

New England Aster is recommended for the following landscape applications;

- Mass Planting
- General Garden Use
- Naturalizing And Woodland Gardens



New England Aster flowers
Photo courtesy of NetPS Plant Finder

Bachman's™

Growing Since 1885

Planting & Growing

New England Aster will grow to be about 3 feet tall at maturity, with a spread of 24 inches. It tends to be leggy, with a typical clearance of 1 foot from the ground, and should be underplanted with lower-growing perennials. The flower stalks can be weak and so it may require staking in exposed sites or excessively rich soils. It grows at a medium rate, and under ideal conditions can be expected to live for approximately 10 years.

This plant does best in full sun to partial shade. It prefers to grow in average to moist conditions, and shouldn't be allowed to dry out. It is not particular as to soil type or pH. It is somewhat tolerant of urban pollution. This species is native to parts of North America. It can be propagated by division.